

**CLINICAL COMPONENT FOR THE DOMICILIARY OXYGEN SERVICE FOR
CHILDREN IN ENGLAND AND WALES**

Dr Ian M Balfour-Lynn

Consultant in Paediatric Respiratory Medicine
Royal Brompton Hospital,
Sydney Street,
London SW3 6NP

Tel: 0207 351 8509

Fax: 0207 351 8763

Email: i.balfourlynn@imperial.co.uk

Dr. Rob A Primhak

Senior Lecturer in Respiratory Paediatrics
Sheffield Children's Hospital
Western Bank
Sheffield S10 2TH

Tel: 0114 271 7585

Fax: 0114 2730522

Email: r.a.primhak@sheffield.ac.uk

Dr Ben NJ Shaw

Consultant in Respiratory & Neonatal Paediatrics
Liverpool Women's Hospital,
Crown Street,
Liverpool L8 7SS

Tel: 0151 708 9988

Fax: 0151 702 4313

Email: ben.shaw@lwh-tr.nwest.nhs.uk

1. Introduction

A separate section is required for paediatric domiciliary oxygen therapy as there are many differences from the adult recommendations. These were initially outlined in a College Report from the Royal College of Physicians Working Group 1992, and although the main document is no longer available, the summary can be obtained [1]. The revised 1999 College Report [2] included a paediatric section, and much of this review has been based on that, together with relevant updates.

It soon became clear that there is a lack of consensus about fundamental paediatric issues such as the ideal target oxygen saturation. This is critical as it determines who needs home oxygen, how much should be given and when it is no longer necessary. Lack of agreement and enormous variety of practice in babies with chronic lung disease has been highlighted by surveys from North America on initiation [3], and from UK and Ireland on weaning, of home oxygen [4]. Although this review attempts to offer guidance to clinicians and service providers, there is a lack of randomised controlled trial evidence to inform much of the debate.

1.2 Principal differences from adult therapy

- Most of the clinical conditions in the infant age group are exclusive and not seen in adults, although sometimes there is overlap between older children and young adults.
- The prognosis in infancy is usually good; many children only need oxygen for a limited period, although this may be for some years.
- Growth and neurodevelopment are important considerations.
- Assessment is different, particularly as regards difficulty of arterial blood sampling.
- Specific equipment is required to allow for low oxygen flows.
- Almost all children receiving long term oxygen therapy (LTOT) also require ambulatory oxygen therapy.
- Many children require LTOT overnight only especially during the weaning stages; this is less than the 15 hours that forms part of the adult LTOT definition.
- All children require supervision from a parent / carer.
- Provision of oxygen may be necessary at school for school-age children.

2. Long Term Oxygen Therapy (LTOT)

2.1 Definition

LTOT refers to the provision of oxygen therapy for continuous use at home for patients with chronic hypoxaemia (due to any cause) in order to maintain oxygen saturation (SaO_2) at or above 92% (depending on the type of oximeter) or arterial PaO_2 above 8kPa (see below). It may be required 24 hours per day or during periods of sleep only. Any child likely to require continuous or intermittent oxygen therapy for more than 2-3 weeks should be considered for discharge home on LTOT. The principal aims of LTOT are to prevent harm from chronic hypoxaemia and to improve any relevant symptoms. It is important to be clear about the aims for an individual child, as this will affect decisions about target oxygenation (hence titration and weaning of flow rates).

2.2 Principal associated conditions

The principal associated conditions are outlined below. The list is not exhaustive, and there are a number of other rare causes of chronic hypoxaemia.

- Chronic neonatal lung disease (bronchopulmonary dysplasia)
- Other neonatal lung conditions *e.g.* pulmonary hypoplasia
- Congenital heart disease with pulmonary hypertension
- Pulmonary hypertension 2° to pulmonary disease
- Interstitial lung disease
- Obliterative bronchiolitis
- End-stage cystic fibrosis and other causes of severe bronchiectasis
- Obstructive sleep apnoea syndrome & other sleep-related disorders
- Neuromuscular conditions requiring non-invasive ventilation
- Disorders of the chest wall *e.g.* thoracic dystrophy, severe kyphoscoliosis
- Palliative care for symptom relief

Chronic neonatal lung disease (CNLD) is the main indication for LTOT in children. The term bronchopulmonary dysplasia (BPD) has come to signify the need for supplemental oxygen at various postnatal or postmenstrual ages, and the definition has changed with time. A recent study of almost 1000 babies born weighing <1000g showed that although the accuracy was limited, use of supplemental oxygen at 36 weeks postmenstrual age was best at predicting long term pulmonary outcome [5]. Infants with CNLD tend to have been born prematurely and are of very low birth weight. Data from surviving babies born weighing <1500g showed 19% developed CNLD [6]. They have often required prolonged assisted ventilation but continued to need supplemental oxygen once extubated. With increased survival of extremely low birthweight premature infants, it is most likely that the incidence of CNLD will also increase [7], as will the need for LTOT. It has been suggested that use of supplementary oxygen in CNLD reduces mortality from sudden infant death [8], and established that it reduces frequency of intermittent desaturations [8], reduces pulmonary hypertension [9], reduces reversible obstructive lung disease [10], improves growth [11,12], and is likely to be beneficial for neurodevelopment [13]. The effect of oxygen on sleep quality is difficult to interpret, but it seems that desaturations are reduced but at the cost of sleep disruption [14-16]. Although there have been no randomised trials of babies on LTOT, it is suggested that caring for babies on supplementary oxygen at home is preferable to a prolonged hospital stay [17]. It reduces the risk of nosocomial infection (although friends and relatives who are unwell should stay away) and it is felt it is good for parent-child bonding [13]. Finally, it is beneficial in terms of freed resources for neonatal units and reduces the total cost of care for an infant [3,17,18].

Other neonatal lung conditions include pulmonary hypoplasia, congenital pneumonia and meconium aspiration syndrome, but compared to CNLD these cases are rare. Survivors of congenital diaphragmatic hernia repair not uncommonly develop chronic lung disease, mainly due to pulmonary hypoplasia or lung damage resulting from mechanical ventilation. Some require domiciliary oxygen but this is unusual beyond 2 years of age [19].

Congenital heart disease (CHD). It is likely that only those with pulmonary hypertension (with or without Eisenmenger syndrome) need to be at home on long term oxygen. In cyanotic CHD, oxygen has little effect in raising SaO₂ and is not indicated, although the degree of polycythaemia may be reduced [20]. However in some cases with chronic left to

right shunting, irreversible pulmonary vascular disease can develop and cause right to left shunting (Eisenmenger syndrome). In the early stages, the resulting pulmonary hypertension may be responsive to oxygen, so this is indicated whilst the child is waiting for transplantation or for palliation. In a small but important controlled study, 100% oxygen given for a minimum of 12 hours per day for up to 5 years significantly improved survival in children with pulmonary vascular disease too severe to have corrective surgery [21]. However this was not the case in a recent 2-year study of adults with advanced Eisenmenger syndrome where nocturnal oxygen had no impact on survival, exercise capacity or quality of life [22]. Some children with CHD who are awaiting corrective surgery (without Eisenmenger syndrome), and who have raised pulmonary artery pressure which is oxygen responsive, may benefit from LTOT. Decisions for all these cases are made by specialist paediatric cardiologists.

Pulmonary hypertension resulting from pulmonary disease results from chronic hypoxia and considerably worsens the overall prognosis [23]. Acute hypoxia causes smooth muscle contraction in pulmonary arteries and sustained hypoxia can cause adaptation of pulmonary endothelium [23]. LTOT reverses or at least slows the progress of the hypoxic-induced changes to the pulmonary vascular bed and can contribute to improved survival [23]. *Primary pulmonary hypertension* has a very poor prognosis in children (median survival <1 year). Some of the children desaturate during sleep (especially during early morning hours) due to mild hypoventilation which may lead to severe dyspnoea, and the resulting hypoxaemia can be eliminated by supplemental oxygen [20]. The children also need oxygen available for when they have viral upper respiratory tract infections, as some tend to desaturate.

Interstitial lung disease represents a spectrum of rare conditions (*e.g.* chronic pneumonitis of infancy, non-specific interstitial pneumonitis, desquamative interstitial pneumonitis, immunodeficiency) in which oxygen exchange is impaired. Many of the children require LTOT along with drug therapy and the outlook is variable. *Obliterative bronchiolitis* leads to severe obstructive lung disease, and whilst it may occur post-viral infection (*e.g.* adenovirus), the cause is often unknown. There is no specific therapy, many of the children require LTOT, and again the outlook is variable.

Cystic fibrosis (CF). As therapy improves, there are less children with CF who require LTOT (estimated to be 1-2% paediatric CF patients). There is surprisingly little evidence for the benefit of LTOT in CF, and whilst it led to an improvement in school or work attendance, there was no effect on mortality rate, frequency of hospitalisation or disease progression [24]. It is therefore recommended that LTOT is reserved for those CF patients who obtain symptomatic relief [25]. This is similar to the indications in *palliative care* where some children feel better for having supplemental oxygen. There are other causes of *bronchiectasis* in children (although in approximately 50% cases no underlying cause is found), and occasionally LTOT is necessary for those with severe disease.

Obstructive sleep apnoea syndrome may require non-invasive ventilation (NIV) if the obstruction can not be relieved. Occasionally supplemental oxygen alone is used if the child does not tolerate the facemask ventilation (for example some children with Down's syndrome). Oxygen does not suppress the ventilatory drive in the majority of children with OSAS but PaCO₂ should be monitored [26]. It is important that children with sickle cell disease and upper airway obstruction do not become hypoxaemic as it can lead to debilitating episodes of sickling [27]. As the incidence of childhood obesity continues to increase, OSAS is likely to become a more frequent problem.

Chronic hypoventilation. There are a number of patients who require long term NIV because of chronic hypoventilation (to control hypercapnia as well as hypoxaemia). Oxygen alone is inadequate for most children with chronic hypoventilation. A UK survey in 1997 estimated the number of children receiving NIV to be 141 of which 93 (65%) were at home [28]. The incidence was shown to have increased over the previous decade and it is likely that this will continue to increase in the future. The children had a variety of conditions, principally neuromuscular disorders (46%), congenital central hypoventilation syndrome (13%), spinal injury (12%), craniofacial syndromes (7%) and bronchopulmonary dysplasia (4%). There were a variety of other less frequent causes (18%). In 35% cases they had access to supplemental oxygen at home as well (personal communication E. Jardine). Core guidelines have suggested children on home ventilation should have a stable oxygen requirement with an FiO_2 of usually less than 40% [29].

2.3 Assessment in infancy

Suitability for domiciliary oxygen therapy should be assessed by a specialist with appropriate experience in the care of the relevant condition; this is usually either a respiratory paediatrician or neonatologist (but may be a paediatric cardiologist, general paediatrician or palliative care specialist). The family must also be assessed as competent to manage home oxygen therapy and be able to cope with all aspects of the baby's care.

In adults, measurement of arterial PaO_2 is considered critical but this is not always possible or practical in infants. In paediatric practice, oxygen saturation measured by pulse oximetry remains the main form of assessment, as results of an arterial stab from a crying (hypoxic) child is unreliable, whilst capillary PaO_2 does not necessarily correlate with arterial PaO_2 [30]. One-off measurements are insufficient, and before discharge, saturations must be measured continuously for at least 6-12 hours, to include periods of sleep, wakefulness and activity/feeding. It is important to include these periods as infants have an increased oxygen requirement during activity and infants with CNLD may develop feeding-related hypoxaemia [31]. In addition, some children may only show evidence of nocturnal hypoxaemia, without daytime hypoxaemia. An instrument should be used that has been validated in infants, and it has been suggested that pulse oximeter readings should be verified by an arterial gas measurement at the start of the study; however this is usually impractical in children and we do not feel it is essential. It must still be remembered however, that correlation between SaO_2 and arterial PaO_2 is such that at 94% saturation the PaO_2 may vary from 9-17 kPa [32]. Although, correlation is particularly poor in the saturation range of 85-90% [33], this is not critical as all infants with $\text{SaO}_2 < 90\%$ should receive LTOT anyway.

Most of the work in infancy relates to CNLD and there is only limited evidence to recommend a minimally acceptable level of oxygenation. The normal SaO_2 is around 96% and supplemental oxygen should be considered for infants who cannot maintain SaO_2 at 93% or above when asleep or quietly awake [8]. Oxygen therapy should be given to achieve a SaO_2 above 92% and recommendations have suggested keeping it at 95% or above [8]. However a recent randomised controlled study (the "BOOST" trial) in 358 premature infants still requiring oxygen at 32 weeks postmenstrual age, showed that maintaining SaO_2 at 95-98% had no advantage over 91-94% in terms of growth and neurodevelopment assessed at 1 year of age (using N-3000 Nellcor oximeters) [34]. The study also found that the group with the higher target oxygen level had an excess of deaths from pulmonary causes, albeit not statistically significant. This is in keeping with another study (the "STOP-ROP" trial on retinopathy of prematurity), in which 649 preterm

infants were randomly assigned to different target oxygen levels (89-94% vs 96-99%) for at least 2 weeks [35]. They used Ohmeda 3470 oximeters that are calibrated to display a SaO₂ lower by 1.6% saturation points compared with other commercial oximeters. They found an increased rate of adverse pulmonary sequelae (pneumonia and exacerbations of CNLD), although not deaths, in the high saturation group when assessed at 3 months after the due date of the infant (13.2 % vs 8.5%). The high saturation group also had more infants still requiring supplemental oxygen at 3 months (47% vs 37%). Oxygen toxicity, particularly in premature infants can inhibit lung healing and contribute to ongoing lung injury [36], so it would seem that therapy should aim to keep SaO₂ at 92-94% in infants, particularly during the preterm period, with no more than 5% of time spent below 90% saturation.

2.4 Assessment in older children

For older children, adult criteria for oxygen prescription may be acceptable, though specific studies on upper and lower limits of PaO₂ for LTOT are lacking. It is important to include an entire night (with at least 4 hours uninterrupted sleep) in case the child only develops nocturnal hypoxaemia. Data for normal overnight SaO₂ measured by pulse oximetry exists, and a recent study in primary school children (mean age 9.3 years) showed that the average SaO₂ was approximately 98% and baseline SaO₂ <97% was uncommon [37]. Furthermore while intermittent desaturations by 4% or more were frequent, the SaO₂ rarely fell to 90% or below. Overnight assessment is important as nocturnal hypoxaemia usually precedes daytime desaturation in children with cystic fibrosis. In children with hypoventilation, such as in neuromuscular disease, it is important to check the CO₂ status, as in the presence of hypercapnia, NIV will be necessary rather than LTOT alone. Measurement of overnight transcutaneous or end-tidal CO₂ is ideal, although a capillary CO₂ taken just as the child wakes up may be useful if the former are not available. Full polysomnography is not usually required.

2.5 Discharge criteria

A multidisciplinary meeting should ensure discharge planning is achieved properly. Extensive collaboration is required between the parents and multidisciplinary team, and in some cases it may be appropriate to include a representative of the service contractor in discharge planning to ensure carers are comfortable with the arrangements. The goal is to select the infants and families most likely to cope at home and to get the timing of discharge right [38].

- Oxygen requirement must be stable with mean SaO₂ of 93% or above, without frequent episodes of desaturation. The SaO₂ should not fall below 90% for more than 5% of the artefact-free recording period.
- Babies should be able to cope with short periods in air without being at risk of rapid deterioration in case their nasal cannulae become dislodged. Generally this will apply to those requiring <0.5 l/min, but this does not mean that some babies with a higher oxygen requirement cannot be on home oxygen.
- No other clinical conditions precluding discharge and the child is medically stable with satisfactory growth. No apnoeic episodes for at least 2 weeks.
- Immunisations up to date. Palivizumab (Synagis) may be considered at the appropriate time of year for infants with CNLD requiring home oxygen.
- Parents are willing and felt to be capable of taking the baby home whilst still on oxygen.

- Home conditions must be satisfactory and preferably a telephone is installed (or use of a mobile phone available). A visit from a member of the home care team is required before discharge.
- Parents are trained and have written information on the use of home oxygen and also cardiopulmonary resuscitation in the case of infants. Vigilance for an empty oxygen supply, dislodged cannulae or blocked valve is critical. A structured education program can be useful [40].
- Advice given about no smoking in the home – it needs to be strongly discouraged but help to do so must be offered.
- Advice must also be given about open flames *e.g.* birthday cake candles [39].
- Older children also need to have training on how to use their oxygen equipment.
- Parents must be advised about travel with cylinders and inform their home and car insurers.
- Appropriate support must be in place *e.g.* community nursing, nurse specialists, health visitor, social worker etc.
- Communication with general practitioner has taken place, and roles are clarified for delivering clinical care.
- Parents must have a list of telephone numbers for advice and emergency help including equipment breakdown.
- Arrangements are in place for open access to the local paediatric unit.

2.6 The prescription

A similar prescription form to the adult one will be used, and will include specific information that may be needed by the contractor in the case of paediatric needs. Hospital specialists (most often respiratory paediatricians or neonatologists) will initiate most oxygen therapy prescriptions for children. As with adult prescriptions, this will need to be copied to the service contractor to ensure delivery of the required service as soon as possible, as well as to the child's general practitioner, and to the Primary Care Trust managing the local domiciliary oxygen service contract.

2.7 Domiciliary Oxygen Record (DOR) forms

Details of the assessment, oxygen therapy prescribed, and information recorded during follow-up visits should be recorded in the child's DOR form. This should also include details of any community and/or general paediatricians involved in a shared-care arrangement with the tertiary centre.

2.8 Equipment

Much of the information on LTOT equipment for adults applies to children but there are some specific issues:

- Oxygen concentrators are usually the preferred devices with back up cylinders for breakdown, and portable cylinders (preferably lightweight) for ambulatory use.
- Oxygen concentrators need two outlets, one in the child's bedroom and one in the main living room area.

- Oxygen cylinders may be more appropriate if initial flow rates are lower than 0.3 l/min and the anticipated duration of oxygen therapy is less than 3 months. A back up cylinder must be available for those on continuous oxygen. Particular attention must be paid to safety and securing of cylinders in the presence of young children.
- Low flow meter (0.1-1 l/min) must be available for infants and very young children.
- Humidification system is often required for those on flow rates above 1 l/min for nasal comfort. Cold bubble humidifiers may be used for this purpose, but they only achieve 40% relative humidity, and are inadequate for direct airway humidification (*e.g.* via a tracheostomy). Heated humidification is less convenient for domestic use, and only effective at flow rates of 4 l/min or higher delivered by a concentrator, as water can block the tubing at lower flow rates.
- Appropriately-sized soft twin-prong nasal cannulae (small children rarely tolerate >2 l/min by nasal cannulae), facemask and non-kinking extension tubing must be provided by the hospital. Stomahesive (or equivalent) should also be provided to protect the skin in those using nasal cannulae.
- Ambulatory equipment must be available as part of the oxygen delivery systems unless oxygen is only required at night. This must be lightweight so that older children can handle it themselves, and for infants it must fit on to a pram/pushchair. Parents need advice on the type of pram to buy *i.e.* one with a metal basket underneath that is safe and strong enough to hold the cylinder.
- Children in wheelchairs need to have a cylinder fitting provided by their wheelchair service to maintain safety.

Oxygen saturation monitor (pulse oximeter) – The issue of whether parents should be provided with their own saturation monitor has been an area of debate. If a child requires continuous monitoring then it is unlikely they are ready for hospital discharge. There is no evidence that provision of oximeters improves the outcome of babies on home oxygen, and in practice may lead to excessive adjustments of the flow rate by the carers [41]. It may give false reassurance, and oxygen saturation is only one aspect of the baby's respiratory status. Some parents, however, request one for spot measurements as a guide to when oxygen needs increasing, during for example, a cold. Nevertheless our recommendation would be that the baby should be seen in these circumstances, usually by the home care nurse, who can then make a full assessment. In addition, the use as an overnight 'alarm' is unsatisfactory due to the number of false alarms mostly from movement artefact. The American Thoracic Society has supported provision of oximeters to parents, but their rationale seems to be mostly cost-based in terms of reducing hospital and office visits [42].

2.9 Patient follow-up arrangements

Arrangements for follow-up after discharge should be co-ordinated by the hospital specialist who has initiated domiciliary oxygen. Liaison must take place between the specialist (usually a respiratory paediatrician or neonatologist), local and/or community paediatrician, general practitioner, nurse specialists (usually based in tertiary respiratory or neonatal units), community paediatric nursing service, occupational therapist and health visitor. A study in Oxford on 55 babies with CNLD at home with LTOT looked at healthcare use in 31 of the babies [17]. They found that they received a median of 43 visits per baby (range 8-173) from a paediatric community nurse, and these lasted a median of 45 minutes; 83% saw a health visitor with a median of 12 visits (range 2-82) and they were for a median of 30 minutes; 83% saw a GP with a median of 6 visits (range 1-140) and they were for a median of 10 minutes. A

lesser proportion also saw a hospital consultant (they almost all had hospital clinic visits however on a median 4 occasions), social worker, physiotherapist and speech therapist.

The oxygen requirement is likely to change with time, and in the case of infants with CNLD it should gradually reduce. Regular home monitoring using pulse oximetry is necessary and should be set up by community/specialist nurses. A visit within the first 24 hours is important to allay parental anxiety. The first formal saturation monitoring should take place within a week, subsequent recordings should occur as clinically indicated, but rarely less often than every 3-4 weeks [41]. Recorded data can then be discussed with the supervising consultant although protocols can be in place for experienced nursing teams to initiate changes. The length of time infants with CNLD remain on LTOT varies but is usually 6-12 months, although some require it for several years.

Infants with CNLD who require home oxygen have more frequent and longer hospital admissions, and more clinic attendances than those sent home without oxygen [43]. This is particularly the case if they had been hospitalised in the first 2 years of life with RSV infection, which had a significant impact on cost of care [44]. In the Oxford study, 41% of the babies required readmission, on a median of 1 occasion (range 0-10), staying for a median 9 days (range 1-64) [17]. The children should be seen regularly by the hospital specialist in clinic in order to monitor the underlying condition as well as growth and neurodevelopment. There must be direct access for the child to be admitted to hospital in the case of any emergency or acute deterioration in the condition, and the parents must have the telephone numbers of the team. Even a simple viral upper respiratory tract infection in a young infant on LTOT may necessitate admission, and winter (particularly the first) can be an anxious time for parents. If the carers believe the child requires an increase in oxygen, they can turn up the flow rate but must then seek advice from the home care team.

It is recommended that influenza immunisation should be given to children with chronic respiratory disease requiring home oxygen, although the vaccine is not licensed for those aged under 6 months. In these cases it is worthwhile immunising household contacts and other caregivers to give some protection to the child [45].

Even when the child comes off oxygen, support must continue as the children may sometimes relapse and require further periods of oxygen therapy after an apparently complete recovery. This is usually related to intercurrent infection.

2.10 When can LTOT be weaned?

The issue of weaning off oxygen principally applies to infants with CNLD and some other neonatal lung conditions. In children with congenital heart disease, only those having successful corrective surgery or transplantation will no longer require LTOT. Some of the young children with interstitial lung disease or obliterative bronchiolitis may improve sufficiently to lose their oxygen dependency. The older children with progressive lung disease such as cystic fibrosis or neuromuscular conditions usually continue to require oxygen for the remainder of their lives. It is not easy to counsel parents as to how long LTOT will be needed. Although capillary blood PaCO₂ measured near term correlated with length of oxygen dependency for a group of babies with CNLD, it was impossible to predict for an individual [46]. However in general those with a higher PaCO₂ are more likely to require oxygen for longer. The length of time infants with CNLD remain on LTOT varies but is usually under 12 months, although some require it for a number of years [47-49]. Persisting symptoms or failure to progress warrants review to rule

out conditions such as tracheo-bronchomalacia, large airway stenosis or granuloma formation, gastro-oesophageal reflux, recurrent aspiration, or unsuspected congenital cardiac disease.

There is enormous variety of practice with regard to criteria used by paediatricians over when to discontinue LTOT in children with CNLD, which reflects lack of evidence on which to base guidelines [4]. Regular reassessment (usually monthly if the child is stable) in the home using a pulse oximeter with a data recording facility is necessary. During periods of weaning or withdrawal of oxygen, more frequent monitoring will be needed. It is not normally necessary for the child to have an oxygen saturation monitor kept in the home, as one can be provided for intermittent monitoring. Recording should include daytime activity, feeding and sleep. Short term awake SaO₂ measurements do not predict prolonged sleeping SaO₂ [50]. The same target saturations used to decide supplementation are used for weaning purposes (92-94%). Saturation targets in the physiological range (95-98%) may be felt to be desirable in infants with pulmonary hypertension. In a study using 2-hour room air challenges, most infants reached their lowest saturations within 40 minutes of discontinuing oxygen and a level of 92% or above best predicted readiness for weaning judged by 6 month follow-up [51]. Furthermore, infants requiring an oxygen flow rate of 0.02 l/kg/min were most likely to be successfully weaned [51].

Some units wean infants from continuous low flow oxygen to night-time and naps only, whilst others maintain continuous oxygen until the child has no requirement at all; there is no evidence to recommend which approach is best. When the oxygen requirement is minimal, the children should have supervised weaning into air with continuous monitoring (that will include active periods and sleep). The babies tend to be weaned down to an oxygen flow rate of 0.1 l/min using a low flow meter, and from that level they can usually be weaned straight to air. Although very (or ultra) low flow meters exist (0.025-0.2 l/min), so that in theory the flow can be reduced further before weaning to air, it is usually unnecessary, and in addition there is a concern that some carers may become confused by the decimal points. Weaning is preferably done at home as it minimises the chances of nosocomial infection, although may sometimes need a brief hospital admission. It is usually prudent to ensure the child has coped with at least one viral upper respiratory tract infection without problems before the equipment is removed from the home, and it should be left there for a few months, especially in the winter.

3.1 Ambulatory oxygen therapy

All children on LTOT require facilities for portable AOT unless they only use night-time oxygen. This particularly applies to the infant age group (who spend periods in the day sleeping) as parents will need to be able to take the baby out the home in order to lead a normal life as possible (shopping, taking other children to school etc.). It is essential that appropriate equipment is available, for example oxygen cylinders that fit on a pram. The duration of AOT will usually increase with older children who are rarely 'house-bound', but a separate source should be provided at school.

3.2 Oxygen in the school

There is a small but important demand for oxygen therapy to be available in schools. Liaison is required between the specialist paediatric respiratory team and the education health services, and this is usually co-ordinated by community paediatric services. The following need particular consideration:

- Oxygen delivery equipment must be lightweight and easy for the child to handle and adjust.
- Safety devices must be in place for stabilising oxygen cylinders or other equipment.
- Insurance cover must be obtained by the school for staff and premises.
- Adequate technical back up must be available.
- School staff must be trained in the use of oxygen therapy.
- School staff must have easily identified healthcare contacts.
- Provision must be made for ambulatory oxygen for the journey to and from school.

4. Short burst oxygen therapy

There are no indications specific to children. In any case, they should be supervised in hospital if they require acute use of short-term oxygen, for example during severe acute asthma.

Acknowledgements

Thanks to Beverly Hicks (occupational therapist Royal Brompton Hospital) and Alison Bedford-Russell (consultant neonatologist Chelsea & Westminster Hospital) for advice over aspects of this document.

5. References

1. Silverman M. Domiciliary oxygen therapy for children. Summary of a report of a Working Group of the Committee on Thoracic Medicine of the Royal College of Physicians. *J Roy Coll Physicians Lond* 1992;26:125-27.
2. Domiciliary oxygen therapy services. Clinical Guidelines and Advice for Prescribers. Report of a working party of the Royal College of Physicians. London 1999.
3. Ellsbury DL, Acarregui MJ, McGuinness GA, Eastman DL, Klein JM. Controversy surrounding the use of home oxygen for premature infants with bronchopulmonary dysplasia. *J Perinatol* 2004;24:36-40.
4. Solis A, Harrison G, Shaw BN. Assessing oxygen requirement after discharge in chronic lung disease: a survey of current practice. *Eur J Pediatr* 2002;161:428-30.
5. Davis PG, Thorpe K, Roberts R, Schmidt B, Doyle LW, Kirplani H. Evaluating "old" definitions for the "new" bronchopulmonary dysplasia. *J Pediatr* 2002;140:555-60.
6. Stevenson DK, Wright LL, Lemons JA, Oh W, Korones SB, Papile LA, Bauer CR, Stoll BJ, Tyson JE, Shankaran S, Fanaroff AA, Donovan EF, Ehrenkranz RA, Verter J. Very low birth weight outcomes of the National Institute of Child Health and Human Development Neonatal Research Network, January 1993 through December 1994. *Am J Obstet Gynecol* 1998;179:1632-9.
7. Hack M, Fanaroff AA. Outcomes of children of extremely low birthweight and gestational age in the 1990s. *Semin Neonatol* 2000;5:89-106.
8. Poets CF. When do infants need additional inspired oxygen? A review of the current literature. *Pediatr Pulmonol* 1998;26:424-28.
9. Abman SH, Wolfe RR, Accurso FJ, Koops BL, Bowman CM, Wiggins JW Jr. Pulmonary vascular response to oxygen in infants with severe bronchopulmonary dysplasia. *Pediatrics* 1985;75:80-4.
10. Tay-Uyboco JS, Kwiatkowski K, Cates DB, Kavanagh L, Rigatto H. Hypoxic airway constriction in infants of very low birth weight recovering from moderate to severe bronchopulmonary dysplasia. *J Pediatr* 1989;115:456-9.
11. Groothuis JR, Rosenberg AA. Home oxygen promotes weight gain in infants with bronchopulmonary dysplasia. *Am J Dis Child* 1987;141:992-95.
12. Hudak BB, Allen MC, Hudak ML, Loughlin GM. Home oxygen therapy for chronic lung disease in extremely low-birth-weight infants. *Am J Dis Child* 1989;143:357-60.
13. Kotecha S, Allen J. Oxygen therapy for infants with chronic lung disease. *Arch Dis Child Fetal Neonatal Ed* 2002;87:F11-14.
14. Harris MA, Sullivan CE. Sleep pattern and supplementary oxygen requirements in infants with chronic neonatal lung disease. *Lancet* 1995;345:831-32.
15. Fitzgerald D, Van Asperen P, Leslie G, Arnold J, Sullivan C. Higher SaO₂ in chronic neonatal lung disease: does it improve sleep? *Pediatr Pulmonol* 1998;26:235-40.
16. Simakajornboon N, Beckerman RC, Mack C, Sharon D, Gozal D. Effect of supplemental oxygen on sleep architecture and cardiorespiratory events in preterm infants. *Pediatrics* 2002;110:884-88.

17. Hallam L, Rudbeck B, Bradley M. Resource use and costs of caring for oxygen-dependent children: a comparison of hospital and home-based care. *J Neonatal Nursing* 1996;2:25-30.
18. Greenough A, Alexander J, Burgess S, Chetcuti PAJ, Cox S, Lenney W, Turnbull F, Shaw NJ, Woods A, Boorman J, Coles S, Turner J. High versus restricted use of home oxygen therapy, health care utilisation and the cost of care in chronic lung disease. *Eur J Ped* 2004;163:292-296.
19. Jaillard SM, Pierrat V, Dubois A, Truffert P, Lequien P, Wurtz AJ, Storme L. Outcome at 2 years of infants with congenital diaphragmatic hernia: a population-based study. *Ann Thorac Surg* 2003;75:250-56.
20. Widlitz A, Barst RJ. Pulmonary arterial hypertension in children. *Eur Resp J* 2003;21:155-76.
21. Bowyer JJ, Busst CM, Denison DM, Shinebourne EA. Effect of long term oxygen treatment at home in children with pulmonary vascular disease. *Br Heart J* 1986;55:385-90.
22. Sandoval J, Aguirre JS, Pulido T, Martinez-Guerra ML, Santos E, Alvarado P, Rosas M, Bautista E. Nocturnal oxygen therapy in patients with Eisenmenger syndrome. *Am J Crit Care Med* 2001;164:1682-87.
23. Higenbottam T, Cremona G. Acute and chronic hypoxic pulmonary hypertension. *Eur Resp J* 1993;6:1207-12.
24. Zinman R, Corey M, Coates AL, Canny GJ, Connolly J, Levison H, Beaudry PH. Nocturnal home oxygen in the treatment of hypoxemic cystic fibrosis patients. *J Pediatr* 1989;114:368-77.
25. Schidlow DV, Taussig LM, Knowles MR. Cystic Fibrosis Foundation consensus conference report on pulmonary complications of cystic fibrosis. *Pediatr Pulmonol* 1993;15:187-98.
26. Marcus CL, Carroll JL, Bamford O, Pyzik P, Loughlin GM. Supplemental oxygen during sleep in children with sleep-disordered breathing. *Am J Respir Crit Care Med* 1995;152:1297-301.
27. Blaisdell CJ. Sickle cell disease and breathing during sleep. *Lung Biology in Health & Disease* 2000;147:755-63.
28. Jardine E, O'Toole M, Paton JY, Wallis C. Current status of long term ventilation in children in the United Kingdom: questionnaire survey. *BMJ* 1999;318:295-99.
29. Jardine E, Wallis C. Core guidelines for the discharge home of the child on long term assisted ventilation in children in the United Kingdom. *Thorax* 1998;53:762-67.
30. Yildizdaş D, Yapicioğlu H, Yilmaz HL, Sertdemir Y. Correlation of simultaneously obtained capillary, venous, and arterial blood gases of patients in a paediatric intensive care unit. *Arch Dis Child*. 2004;89:176-80.
31. Singer L, Martin RJ, Hawkins SW, Benson-Szekely LJ, Yamashita TS, Carlo WA. Oxygen desaturation complicates feeding in infants with bronchopulmonary dysplasia after discharge. *Pediatrics* 1992;90:380-4.
32. Wasunna A, Whitelaw AG. Pulse oximetry in preterm infants. *Arch Dis Child* 1987;62:957-58.
33. Roberts CM, Bugler JR, Melchor R, Hetzel MR, Spiro SG. Value of pulse oximetry in screening for long-term oxygen therapy requirement. *Eur Resp J* 1993;6:559-62.

34. Askie LM, Henderson-Smart DJ, Irwig L, Simpson JM. Oxygen-saturation targets and outcomes in extremely preterm infants. *N Engl J Med* 2003;349:959-67.
35. Supplemental Therapeutic Oxygen for Prethreshold Retinopathy Of Prematurity (STOP-ROP), a randomized, controlled trial. I: primary outcomes. *Pediatrics* 2000;105:295-310.
36. Weinberger B, Laskin DL, Heck DE, Laskin JD. Oxygen toxicity in premature infants. *Toxicol Appl Pharmacol* 2002;181:60-7.
37. Urschitz MS, Wolff J, Von Einem V, Urschitz-Duprat PM, Schlaud M, Poets CF. Reference values for nocturnal home pulse oximetry during sleep in primary school children. *Chest* 2003;123:96-101.
38. Gracey K, Talbot D, Lankford R, Dodge P. The changing face of bronchopulmonary dysplasia: Part 2. Discharging an infant home on oxygen. *Adv Neonatal Care*. 2003;3:88-98.
39. Laubscher B. Home oxygen therapy: beware of birthday cakes. *Arch Dis Child* 2003;88:1125.
40. Brown KA, Sauve RS. Evaluation of a caregiver education program: home oxygen therapy for infants. *J Obstet Gynecol Neonatal Nurs* 1994;23:429-35.
41. Primhak RA. Discharge and aftercare in chronic lung disease of the newborn. *Semin Neonatol* 2003;8:117-26.
42. American Thoracic Society. Statement on the care of the child with chronic lung disease of infancy and childhood. *Am J Respir Crit Care Med* 2003;168:356-396.
43. Greenough A, Alexander J, Burgess S, Chetcuti PA, Cox S, Lenney W, Turnbull F, Shaw NJ, Woods A, Boorman J, Coles S, Turner J. Home oxygen status and rehospitalisation and primary care requirements of infants with chronic lung disease. *Arch Dis Child* 2002;86:40-3.
44. Greenough A, Alexander J, Burgess S, Bytham J, Chetcuti PAJ, Hagan J, Lenney W, Melville S, Shaw NJ, Boorman J, Coles S, Turner J. Health care utilisation of prematurely born, preschool children related to hospitalisation for RSV infection. *Arch Dis Child* 2004;89:673-678.
45. Munoz FM. Influenza virus infection in infancy and early childhood. *Paediatr Respir Rev* 2003;4:99-104.
46. Victor S, Shaw B. Carbon dioxide levels do not predict duration of home oxygen requirement: a retrospective study. *J Perinat Med* 2002;30:333-35.
47. Sauve RS, McMillan DD, Mitchell I, Creighton D, Hindle NW, Young L. Home oxygen therapy. Outcome of infants discharged from NICU on continuous treatment. *Clin Pediatr* 1989;28:113-18.
48. Baraldi E, Carra S, Vencato F, Filippone M, Trevisanuto D, Milanese O, Pinello M, Zanardo V, Zacchello F. Home oxygen therapy in infants with bronchopulmonary dysplasia: a prospective study. *Eur J Pediatr* 1997;156:878-82.
49. Hazinski TA. Bronchopulmonary dysplasia. In: Chernick V, Boat TF, eds. *Kendig's disorders of the respiratory tract in children*. Philadelphia, PA, USA: WB Saunders, 2000:364-85.
50. Moyer-Mileur LJ, Nielson DW, Pfeffer KD, Witte MK, Chapman DL. Eliminating sleep-associated hypoxemia improves growth in infants with bronchopulmonary dysplasia. *Pediatrics* 1996;98:779-83.

51. Simoes EA, Rosenberg AA, King SJ, Groothuis JR. Room air challenge: prediction for successful weaning of oxygen-dependent infants. *J Perinatol* 1997;17:125-29.